

MOTRIN® Professional Dosing Information for Infants & Children

Note: If possible, use weight to dose, otherwise use age. To arrive at the correct dose, weigh your child before giving MOTRIN®. A healthcare professional should be consulted for dosing in children under the age of six months.



Always ask your healthcare professional which product is right for your child.

		Your child's weight and age									
		Age 0-5 mos	6-11 mos	12-23 mos	2-3 yrs	4-5 yrs	6-8 yrs	9-10 yrs	11 yrs		
Weight		6-11 lbs	12-17 lbs	18-23 lbs	24-35 lbs	36-47 lbs	48-59 lbs	60-71 lbs	72-95 lbs		
 <p>Concentrated MOTRIN® Infants' Drops Berry & Berry dye-free flavors Active Ingredient: Ibuprofen 50mg (in each 1.25mL) Use only as directed.</p>		—	1.25mL	1.875mL	—	—	—	—	—		
	 <p>Children's MOTRIN® Suspension Berry, Grape, Tropical Punch, Bubblegum & Berry dye-free flavors Active Ingredient: Ibuprofen 100mg (in each 5mL or 1 tsp) Use only as directed.</p>		—	—	—	1 tsp or 5mL	1 1/2 tsp or 7.5mL	2 tsp or 10mL	2 1/2 tsp or 12.5mL	3 tsp or 15mL	
		 <p>MOTRIN® Junior Strength Easy-to-Swallow Caplets Active Ingredient: Ibuprofen 100mg (in each caplet) Use only as directed.</p>		—	—	—	—	—	2 caplets	2 1/2 caplets	3 caplets
			 <p>MOTRIN® Junior Strength Chewable Caplets Orange & Grape flavors Active Ingredient: Ibuprofen 100mg (in each tablet) Use only as directed.</p>		—	—	—	1 tablet	1 1/2 tablets	2 tablets	2 1/2 tablets

tsp = teaspoon*

Take every 6-8 hours as needed. Do not exceed more than 4 doses in 24 hours.

Important Instructions for Proper Use

- **Read and follow the label instructions on all MOTRIN® products**
 - Take every 6-8 hours as needed
 - Do not exceed more than 4 doses in 24 hours
- **Do NOT administer longer than 10 days, unless directed by a doctor**
- **Ask a doctor or pharmacist before use if the child is taking any other drug containing an NSAID (prescription or non-prescription)**
 - MOTRIN® contains ibuprofen
- **Keep all medicines out of the reach of children**
- ***Use only the dosing device that comes with a specific product**
 - Household items such as measuring spoons are less accurate
- **Do NOT administer adult medicines to children**
- Concentrated MOTRIN® Infants' Drops are more concentrated than Children's MOTRIN® Liquids. The Concentrated Infants' Drops have been specifically designed for use with enclosed dosing device.
- Children's MOTRIN® Liquids are less concentrated than Concentrated MOTRIN® Infants' Drops. The Children's MOTRIN® Liquids have been specifically designed for use with the enclosed measuring cup.
- The OTC dosing: the recommended dose is 7.5mg/kg every 6-8 hours. The recommended daily dose is 30mg/kg.
- **If you have any questions, contact your healthcare professional or call 1-877-895-3665**